



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C07F 9/38, A61K 31/66, C07F 9/576, C07C 215/04, C07D 295/08, C07J 51/00, A61K 9/127, C07F 9/572, 9/58, 9/6561	A1	(11) International Publication Number: WO 99/02539 (43) International Publication Date: 21 January 1999 (21.01.99)
(21) International Application Number: PCT/EP98/04238 (22) International Filing Date: 8 July 1998 (08.07.98) (30) Priority Data: MI97A01628 9 July 1997 (09.07.97) IT (71) Applicants (for all designated States except US): GEANGE LIMITED [IE/IE]; 17 Earlsfort Terrace, Dublin 2 (IE). EURODRUG LTD. [CN/CN]; Java Comm. Centre, Room 1501, 124-134 Java Road, North Point, Hong Kong (CN). LIPOTEC S.A. [ES/ES]; Calle Santa Eulalia 240, E-08902 L'Hospitalet de Llobregat (ES). (72) Inventors; and (75) Inventors/Applicants (for US only): VAN GULIK, Fred [NL/CN]; Eurodrug Ltd., Java Comm. Centre, Room 1501, 124-134 Java Road, North Point - Hong Kong (CN). PARENTE, Antonio [ES/ES]; Lipotec s.a., Calle Santa Eulalia 240, E-08902 L'Hospitalet de Llobregat (ES). (74) Agents: TRUPIANO, Federica et al.; Brevetti Europa S.r.l., Piazza Bernini, 6, I-20133 Milano (IT).		(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, GM, HR, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i>
(54) Title: DIPHOSPHONIC ACID SALTS FOR THE TREATMENT OF OSTEOPOROSIS (57) Abstract Disphosphonates, or their salts between disphosphonic acids and linear, branched substituted and non-substituted, cyclic, hetero-cyclic and aromatic amino-alcohols, and their use in the treatment of osteoporosis, and pharmaceutical compositions that contain at least one diphosphonate as active principle for the treatment of osteoporosis.		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon	KR	Republic of Korea	PL	Poland		
CN	China	KZ	Kazakhstan	PT	Portugal		
CU	Cuba	LC	Saint Lucia	RO	Romania		
CZ	Czech Republic	LI	Liechtenstein	RU	Russian Federation		
DE	Germany	LK	Sri Lanka	SD	Sudan		
DK	Denmark	LR	Liberia	SE	Sweden		
EE	Estonia			SG	Singapore		

DISPHOSPHONIC ACID SALTS FOR THE TREATMENT OF OSTEOPOROSIS

OBJECT OF THE PRESENT INVENTION

5

Object of the present invention are salts of diphosphonic acids or diphosphonates and a procedure for their preparation.

Object of the present invention is also the use of mentioned salts of diphosphonic acids or diphosphonates for treatment of osteoporosis.

10

Object of the present invention are again pharmaceutical compositions which contain, as active principle, the mentioned salts of diphosphonic acids or diphosphonates and the use of these compositions in the treatment of osteoporosis.

15

STATUS OF THE TECHNIQUE

As well known osteoporosis is a disease of the skeleton that is characterised by the disarrangement and by the loss of bone tissue, with increase of skeleton fragility and predisposition to fractures.

As an example, is reported that in US this disease
20 afflicts more than 25 millions of persons and it causes 1.300.000 fractures every year.

The social cost is consequently very high accounting for more than 10 billions of US dollars per year. The same figures are reported in Europe and in Japan.

25

CONFIRMATION COPY

SUBSTITUTE SHEET (RULE 26)

The elderly is the population with the highest risk. Unfortunately the problem is becoming progressively more serious in the view of the progressive grow old of population. A forecast of the fracture incidence in the
5 future sixty years assumes an increase of at least three times. Women are more exposed than men, particularly during the first 5 years after the menopausal period.

At present, from a therapeutic viewpoint, either a mono-therapy with fluorides, as reported by Melmon & Morrelli,
10 Clinical Pharmacology: Basic Principles in Therapeutics, 497 (1992), N. Engl. J. Med. 306,446 (1982) is used, or addition use of combined therapies with calcium, fluoride and oestrogens are reported to retain some advantages as compared to the mono-therapy.

15 Alternatively, another class of compounds prescribed for the treatment of osteoporosis includes disphosphonates; these agents are mostly administered by oral route, but besides having the unsolved inconvenient of a poor absorption (from 1 to 5% of dose), they also develop, as
20 primary adverse events, oesophagitis and gastrointestinal disorders of moderate-severe intensity.

To overcome this inconvenient, several clinical studies have been carried out, in the search of new compounds
25 that combine a high pharmacological activity together

with a better tolerability and increased bioavailability as compared to known available compounds.

For example, in the treatment of urolithiasis and for inhibition of bone resorption, diphosphonic acids and their salts are described in the US patent 4,621,077; namely, the biological activities of mentioned phosphonic acids or their corresponding sodium salts are reported. The degree of absorption, i.e. the bioavailability, of those compounds resulted however very low and for this reason, in order to achieve in the body active concentrations of the active principle, high doses had to be administered with consequent occurrence of gastrointestinal adverse reactions which worsen tolerability.

OBJECTIVES OF THE INVENTION

Objective of the present invention is to make available diphosphonates that have an increased bioavailability when administered by oral route.

Objective of the present invention is also to make available diphosphonates that are more effective in the treatment of osteoporosis while retaining a lower incidence of oesophagus and gastro-intestinal adverse events as compared to those known compounds already available.

Again, objective of the present invention is to make available diphosphonates which can be administered both by oral or intravenous route.


5 Again, objective of the present invention is to make available diphosphonates that display an improved therapeutic index in treatment of osteoporosis as compared to the known drugs at present used in the medical practice.

10 Objective of the present invention is also to make available a procedure for the preparation of diphosphonates.

15 Objective of the present invention is also to make available pharmaceutical formulations, containing at least one diphosphonate as active principle, easy to be administered, well tolerated and able to allow a high therapeutic index.

DESCRIPTION OF THE INVENTION

20 These and other objectives with further advantages which are clarified in the description below, are obtained by diphosphonates or salts of diphosphonic acids and linear, branched, substituted and non-substituted, cyclic, hetero-cyclic, aromatic amino-alcohol derivatives, said diphosphonates having the following general formula:




where $R=R_1$ or $R \neq R_1$ and R and R_1 are chosen among H, OH, Cl, linear or branched, substituted or not substituted alkyl groups, linear or branched, substituted or not substituted alkylidenic chain, or they are chosen among:

10


15

20


25




(II)




(III)




(IV)




(V)



(VI)



(VII)




5 More particularly, according to the present invention, mentioned amino-alcohols are chosen among linear and branched amino-alcohols, and specifically they are chosen among:

10


15

20


25




5




10



15




20




Always according to the present invention, mentioned amino-alcohols are cyclic and hetero cyclic amino-alcohols and are chosen among


25




5



10




15




Again, according to the present invention, mentioned
substituted alkylidenic chain, defined in formula (I), is
chosen equal to:

25







with n within 1 and 6, where $\text{R}_2=\text{R}_3$ or $\text{R}_2\neq\text{R}_3$ and R_2 and R_3 are chosen among H, linear or branched, cyclic, substituted or non-substituted alkyl groups, or R_2 and R_3 represent, together with the nitrogen atom N, an aliphatic or aromatic hetero-cycle.


Always according to the present invention, the substituted alkylidenic chain, defined in formula (I), is also chosen equal to




where n is within 1 and 6 and R_4 is chosen among:



According to the present invention, of particular interest appeared the salt obtained between a diposphonic acid of general formula (I), where $R \neq R_1$, R is chosen equal to OH and R_1 is chosen as substituted linear alkyldenic chain of formula (X), where $n=3$, R_2 and R_3 are equal to H according to the following formula:



or 4-amino-1-hydroxybutane-1,1-biphosphonic acid (alendronic acid) and one branched amino-alcohol of formula



or 2-amino-2-hydroxymethyl-1,3-propanediol (trometamine).

15 Mentioned salts, always according to the present invention, depending upon the preparation procedure, are formed by the residues corresponding to the diphosphonic acid and the amino-alcohol in a variable molar ratio, while they can be advantageously used in the treatment of osteoporosis. Mentioned molar ratios can vary between 1:1
20 and 1:2 (diphosphonic acid : amino-alcohol).

Diphosphonates according to the prior art, retain a non satisfactory pharmacokinetic profile, characterised by a very poor absorption from the gastro-intestinal tract,
25 with a systemic bioavailability ranging between 1 to 5 %

of the oral dose. About 50% of the absorbed dose is then eliminated with the urine in 24 hours, while the dose left is retained by bone tissue and afterwards eliminated very slowly.. The unabsorbed diphosphonate is instead
5 eliminated with the faeces.

This behaviour has to be attributed to the physico-chemical properties of the products, which being highly hydro-soluble hardly penetrate the lipid membranes. Oral doses much higher than those bioavailable are then
10 necessary to achieve and maintain a therapeutic effective concentration at the target site (bone), with consequent increase of the local gastro-intestinal adverse events like nausea and diarrhoea.

Diphosphonates according to the present invention, due to
15 the lower dissociation degree of the salt and to the physico-chemical properties of the amino-alcohol, are characterised by a lower water solubility and an increased lipid solubility as compared to the known diphosphonates (sodium salts).

20 These characteristics confer remarkable advantages in the absorption process trough the biological membranes, thus improving bioavailability.

In addition, diphosphonates of the present invention, did
25 show a promising pharmaco-toxicological profile in


relation to the expected use in the therapy of osteoporosis.

The diphosphonates of the present invention show a good efficacy in stimulating bone growth, this being probably
5 due to the known property of amino-alcohols of subtracting toxic-radicals produced by reactive species.


The diphosphonates of the present invention can be easily administered both by intravenous and oral route. In this
10 latter case, the use of pharmaceutical systems able to control the release of the active principle, are particularly advantageous. These systems include liposoms, niosomes and similar matrices, possibly added appropriate carriers which are of easy use and allow a
15 sustained release of the active principle.

The diphosphonates of the present invention are advantageously prepared by reacting a solution or
suspension of diphosphonic acid of general formula (I) with an amino-alcohol dissolved in a proper solvent; the
20 reaction product (salt) between the diphosphonic acid and the amino-alcohol, obtained by cooling the resulting solution, is then purified according to known techniques.

As a non-limitative example of the present invention, some examples referring to the preparation of the salt
25 obtained by reacting the diphosphonic acid of formula:



and the amino-alcohol of formula:



are hereafter described.

Example 1

15 4-amino-1-hydroxybutane-1,1-biphosphonic acid,
 (alendronic acid, 1g, 4.016 mmoles), was dissolved into 20
 mL of CH_2Cl_2 / methyl alcohol (9:1). 2-amino-2-
 hydroxymethyl-1,3 propandiol, (trometamine, 486 mg, 4.016
 20 mmoles) dissolved in 5 mL of CH_2Cl_2 / methyl alcohol
 (9:1) was added under stirring to the alendronic acid
 solution. The white precipitate formed, was filtered and
 dried under vacuum.

1.263 g of alendronate trometamine salt were obtained,
 25 yield 85%.

Example 2

4-amino-1-hydroxybutane-1, 1-biphosphonic acid,
(alendronic acid, 1g, 4.016 mmoles), was suspended into
20 mL water. 2-amino-2-hydroxymethyl-1, 3 propandiol,
5 (trometamine, 486 mg, 4.016 mmoles) was added, under
stirring, to the aqueous solution of alendronic acid.
Solids rapidly dissolved in water, the suspension becomes
clear, while the resulting solution, having a pH = 4.18,
was frozen and lyophilised to give a white powder.

10 1.480 g of alendronate trometamine salt were obtained,
yield 99%.

Example 3

4-amino-1-hydroxybutane-1, 1-biphosphonic acid,
15 (alendronic acid, 1g, 4.016 mmoles), was suspended into 40
mL water. 2-amino-2-hydroxymethyl-1, 3 propandiol,
(trometamine, 972 mg, 8.032 mmoles) was added, under
stirring, to the aqueous solution of alendronic acid.
Solids rapidly dissolved in water, the suspension becomes
20 clear, while the resulting solution had a pH = 7.14 was
frozen and lyophilised to give a white powder.

1.97 g of alendronate trometamine salt were obtained,
yield 99%.

Example 4

4-amino-1-hydroxybutane-1, 1-biphosphonic acid,
(alendronic acid, 1g, 4.016 mmoles), was suspended into 5
mL water. 2-amino-2-hydroxymethyl-1, 3 propandiol,
trometamine (486 mg, 4.016 mmoles) was added, under
5 stirring, to the aqueous solution of alendronic acid.
Solids rapidly dissolved in water, the suspension became
clear, while the resulting solution had a pH = 4.18. 10
mL of iso-propyl alcohol were added and the resulting
solution was cooled at a temperature of -10 °C overnight.
10 The precipitate formed was filtered, washed twice with
cool iso-propyl alcohol, then was dried under vacuum.

743 mg of alendronate trometamine salt were obtained,
yield 50%.


15 The product obtained according to the procedures reported
in the examples 1, 2 and 4, was analysed by IR and ¹H-NMR
spectroscopy.

The melting point was 180.53 °C.


¹H-NMR (300 MHz), (D₂O) δ=3.615 (s 6H, CH₂O) ; 2.920 (t,
20 2H, N⁺-CH₂ , J=6.5 Hz); 1.900 (m, 2H, NCH₂ CH₂ CH₂); 1.890
(t, 2H, NCH₂ CH₂ CH₂ , J=6.5 Hz).

As an example, a non-limitative example is hereafter
reported, referring to the preparation of a
25 pharmaceutical formulation of the salt obtained by

reacting the diphosphonic acid of formula :



and the amino-alcohol of formula :



Example 5

15 Alendronate trometamine salt formulated in liposome matrix.

- LECINOL LS (liposomes) , 20%

- SALT OF 4-AMINO-1-HYDROXYBUTANE-1, 1-BIPHOSPHONIC ACID,
(ALENDRONIC ACID) AND 2-AMINO-2-HYDROXYMETHYL-1, 3

20 PROPANDIOL, (TROMETAMINE) , 15%

(Ratio between alendronic acid and trometamine \cong 1:1)

- TROMETAMINE BASE; 1%

- CHOLESTEROL; 4%

- PROPYLENGLYCOL; 10%

25

- ETHANOL; 1%

- CARBOPOL ETD 2020; 0,5%

- Na₂EDTA ; 0.15%

- NIPAGIN; 0.2%

5 - PHENONIP (CONSERVATIVE); 0,3%

- CITRIC ACID; 0.2

- BHT; 0.01

- WATER; to 100 %

10 Dealing with the evaluation of the pharmaco-toxicological activity of the salts obtained by reacting diphosphonic acids of general formula (I) and amino-alcohols according to the present invention, biological tests were carried out taking as test compound the product alendronate trometamine obtained as described in the examples above.

15

EFFICACY

A study on a tissue culture (scalp slices of new-born mouse prepared according to the method described by A. Togari, Gen. Pharmacol. 24, 1133, 1993) was performed. At
20 the end of the culture period, the tissue added μ molar concentrations of the title product, was homogenised in saline plus 0.1% Triton X-100.

25

The obtained homogenate was used to determine alkaline phosphatase (as osteoblastic activity index) and N-acetylglycosaminidase (as osteoclastic activity index).

5 The test product starting from the 10 μ molar concentration, was shown to be able to activate in a significant manner the osteoblastic activity (alkaline phosphatase) and to inhibit the osteoclastic activity (N-acetylglycosaminidase).

10 TOXICITY

The compound when administered orally to mice at the dose of 50 mg/kg, which is a multiple (x300 times) of the human therapeutic dose, did not cause the onset of apparent toxic symptoms.

15

BIOAVAILABILITY

A study carried out in rats administered both orally and intravenously the test compound, alendronate trometamine, at the 10 mg/kg dose allowed calculation of a mean
20 absolute bioavailability equal to 8%. Calculation was performed on 0-24 hours urine elimination of alendronic acid according to the method described in J. of Chromatography, 533, 183-193, 1992: "Improved
25 determination of the biphosphonate alendronate in human

plasma and urine by automated pre-column derivatisation and HPLC with fluorescent and electro-chemical detection".

5

10

15


20

25

CLAIMS

1. Diphosphonates or salts of diphosphonic acids and linear, branched, substituted and not substituted, cyclic, heterocyclic, aromatic amino-alcohol derivatives

5 having the following general formula:




10 where $\text{R}=\text{R}_1$ or $\text{R}\neq\text{R}_1$ and R and R_1 are chosen among H, OH, Cl, linear or branched, substituted or not substituted alkyl groups, linear or branched, substituted or not substituted alkylidenic chain, or they are chosen among:

15


20

25

23




(II)




(III)

5




(IV)

10



(V)

15




(VI)

20



(VII)

25



2. Diphosponates according to Claim 1, characterised by the fact that the mentioned amino-alcohols are chosen among linear and branched amino-alcohols, and specifically they are chosen among:


10

15


20

25


5




10




15



20




25




3. Diphosponates according to Claim 1, characterised by the fact that the mentioned amino-alcohols are cyclic and hetero cyclic amino-alcohols and are chosen among


5




10



15




20



25


4. Diphosponates according to Claim 1, characterised by the fact that mentioned linear or branched substituted or non substituted alkylidenic chain , defined in formula (I), is chosen equal to:

5



with n within 1 and 6, where $\text{R}_2=\text{R}_3$ or $\text{R}_2\neq\text{R}_3$ and R_2 and R_3 are chosen among H, linear or branched, cyclic, substituted or not substituted alkyl groups, or R_2 and R_3 represent together with the nitrogen atom N an aliphatic or aromatic hetero-cycle.


5. Diphosponates according to Claim 1. characterised by the fact that mentioned substituted alkylidenic chain, defined in formula (I), is also chosen equal to




where n is within 1 and 6 and R_4 is chosen among:

20


25




(XII)



(XIII)




(XIV)




(XV)

6. Diphosponate according to Claim 1, characterised by the fact that is a salt obtained between a diphosphonic acid of general formula (I), where $R \neq R_1$, R is chosen equal to OH and R_1 is chosen as substituted linear alkyldienic chain of formula (X), where $n=3$, R_2 and R_3 are equal to H according to the following formula:



(XXVI)

or 4-amino-1-hydroxybutane-1,1 bi-phosphonic acid (alendronic acid) and one branched amino-alcohol of formula



or 2-amino-2-hydroxymethyl-1,3-propanediol (trometamine).

7. Process for the preparation of diphosphonates according to claim 1, comprising the following steps:

- mixing of a diphosphonic acid of general formula (I) dissolved or suspended in appropriate solvent, with an amino-alcohol, possibly also dissolved in a proper solvent.
- 15 • Cooling of the mixture obtained and/or possible treatment with a proper solvent, to obtain a diphosphonate or salt of the mentioned diphosphonic acid of general formula (I) and the mentioned amino-alcohol
- 20 • Filtration, washing, possible purification and dryness (lyophilisation) of mentioned product (salt).

8. Use of diphosphonates or their salts between a diphosphonic acid of general formula (I) and amino-

alcohols as in Claim 1, for the treatment of osteoporosis,

5 9. Pharmaceutical compositions for the treatment of osteoporosis comprising at least one diphosphonate or a salt between a diphosphonic acid of general formula (I) and an amino- alcohol as in Claim 1, as active principle.

10 10. Pharmaceutical compositions, according to Claim 9, characterised by the fact that are prepared for parenteral (intravenous) administration.

11. Pharmaceutical compositions, according to Claim 9, characterised by the fact that includes liposomes or niosomes and proper carriers which allow the controlled release of the active principle.

12. Pharmaceutical compositions, according to Claim 9, characterised by the fact that they have the following composition:

20 - LECINOL LS (liposomes) , 20%
- SALT OF 4-AMINO-1-HYDROXYBUTANE-1, 1-BIPHOSPHONIC ACID, (ALENDRONIC ACID) AND 2-AMINO-2-HYDROXYMETHYL-1, 3 PROPANDIOL, (TROMETAMINE) , 15%

(Ratio between alendronic acid and trometamine $\cong 1:1$)

25 - TROMETAMINE BASE; 1%

31

- CHOLESTEROL; 4%
- PROPYLENGLYCOL; 10%
- ETHANOL; 1%
- CARBOPOL ETD 2020; 0,5%
- 5 - Na₂EDTA ; 0.15%
- NIPAGIN; 0.2%
- PHENONIP (CONSERVATIVE); 0,3%
- CITRIC ACID; 0.2
- BHT; 0.01
- 10 - WATER; to 100 %

13. Pharmaceutical composition, according to Claim 12, characterised by the fact that is under liposomal form

14. Diphosphoates according to Claim 1, characterised by the fact that molar ratios between mentioned
- 15 diphosphonic acids and the amino-alcohols are in a molar ratio between 1:1 and 1:2.

20

25

INTERNATIONAL SEARCH REPORT

International Application No

PCT/EP 98/04238

A. CLASSIFICATION OF SUBJECT MATTER

IPC 6	C07F9/38	A61K31/66	C07F9/576	C07C215/04	C07D295/08
	C07J51/00	A61K9/127	C07F9/572	C07F9/58	C07F9/6561

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 C07F C07C C07D C07J A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	CHEMICAL ABSTRACTS, vol. 121, no. 11, 12 September 1994 Columbus, Ohio, US; abstract no. 125254, YONESHIMA N ET AL: "1,3,5(10)-estratriene-3,17.beta.-diol 17-(3,3-diphosphono)propionate for prevention and treatment of osteoporosis" XP002081479 see abstract & JP 06 100576 A (KIRIN BREWERY; JAPAN) 12 April 1994	1-14
Y	GB 2 118 042 A (INSTITUTO GENTILI SPA) 26 October 1983 cited in the application see the whole document --- -/--	1-14

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance
 "E" earlier document but published on or after the international filing date
 "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
 "O" document referring to an oral disclosure, use, exhibition or other means
 "P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
 "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
 "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
 "&" document member of the same patent family

Date of the actual completion of the international search

20 October 1998

Date of mailing of the international search report

05/11/1998

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
 NL - 2280 HV Rijswijk
 Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
 Fax: (+31-70) 340-3016

Authorized officer

Beslier, L

INTERNATIONAL SEARCH REPORT

Int ernational Application No
PCT/EP 98/04238

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No. —
X	CHEMICAL ABSTRACTS, vol. 126, no. 21, 26 May 1997 Columbus, Ohio, US; abstract no. 281009, SAO T ET AL: "Scale dissolving aqueous solutions containing alkanolamine salts" XP002081480 see abstract & JP 09 053193 A (MITSUBISHI HEAVY IND LTD;JAPAN) 25 February 1997 ---	1,2
X	FR 2 142 051 A (HERCULES INC.) 26 January 1973 see example 23; table III ---	1,2
X	EP 0 292 190 A (LION CORP.) 23 November 1988 see page 3, line 44-45 ---	1,2
X	WO 92 00721 A (THE PROCTER & GAMBLE CO.) 23 January 1992 see example 5 ---	1,2
X	US 4 830 779 A (SEIJI MAENO) 16 May 1989 see column 3, line 57-60 ---	1,2
X	GB 1 508 772 A (SHELL INTERNATIONALE RESEARCH MAATSCHAPPIJ BV) 26 April 1978 see examples 36,39,45 ---	1,2
X	US 5 389 261 A (JOHN DALY) 14 February 1995 see claims 2,6,7; example 8 ---	1,2
X	EP 0 781 804 A (KAO CORP.) 2 July 1997 see page 7; example 27 ---	1,2
X	DD 273 650 A (VEB ROBOTRON-ELEKTRONIK RADEBERG) 22 November 1989 see example 4 ---	1,2
X	DD 150 744 A (WOBST, SIGURD) 16 September 1981 see page 8; example 2 -----	1,2

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/EP 98/04238

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
GB 2118042	A	26-10-1983	BE 896453 A	01-08-1983
			CH 661437 A	31-07-1987
			DE 3313049 A	17-11-1983
			FR 2525223 A	21-10-1983
			HK 79390 A	12-10-1990
			JP 1589937 C	30-11-1990
			JP 2013645 B	04-04-1990
			JP 58189193 A	04-11-1983
			LU 84746 A	17-11-1983
			LU 88714 A	23-08-1996
			MX 9203428 A	01-07-1992
			NL 8301324 A, B,	01-11-1983
			SE 463239 B	29-10-1990
			SE 8302130 A	16-10-1983
			US 4621077 A	04-11-1986
FR 2142051	A	26-01-1973	AU 466454 B	20-12-1973
			AU 4342272 A	20-12-1973
			BE 785050 A	18-12-1972
			CA 979318 A	09-12-1975
			GB 1386814 A	12-03-1975
			NL 7208291 A	20-12-1972
			SE 380242 B	03-11-1975
			US 3751372 A	07-08-1973
EP 292190	A	23-11-1988	JP 63286478 A	24-11-1988
			DE 3878734 A	08-04-1993
WO 9200721	A	23-01-1992	AT 130185 T	15-12-1995
			AU 8231191 A	04-02-1992
			CA 2086620 A, C	14-01-1992
			DE 69114708 D	21-12-1995
			DE 69114708 T	02-05-1996
			DK 539480 T	11-03-1996
			EP 0539480 A	05-05-1993
			ES 2079673 T	16-01-1996
			GR 3018102 T	29-02-1996
			IE 71649 B	26-02-1997
			PT 98309 A	29-05-1992
			SK 10193 A	06-10-1993

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/EP 98/04238

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
US 4830779 A	16-05-1989	JP 63165457 A DE 3743928 A	08-07-1988 07-07-1988
GB 1508772 A	26-04-1978	NONE	
US 5389261 A	14-02-1995	CA 2077312 A	02-03-1994
EP 781804 A	02-07-1997	JP 9176427 A CN 1157300 A	08-07-1997 20-08-1997
DD 273650 A	22-11-1989	NONE	
DD 150744 A	16-09-1981	NONE	